

United States Department of Agriculture

Frenchman Watershed



Hydrologic Unit Code 10250005


Natural Resources Conservation Service

Rapid Assessment

Lakewood, Colorado

RWA 10250005

February 2008



The United States Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.)

Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington DC 20250-9410, or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

Introduction

Background Information

The Natural Resources Conservation Service (NRCS) is encouraging the development of rapid watershed assessments in order to increase the speed and efficiency generating information to guide conservation implementation, as well as the speed and efficiency of putting it into the hands of local decision makers.

Rapid watershed assessments provide initial estimates of where conservation investments would best address the concerns of landowners, conservation districts, and other community organizations and stakeholders. These assessments help landowners and local leaders set priorities and determine the best actions to achieve their goals.


Benefits of these Activities

While rapid assessments provide less detail and analysis than full-blown studies and plans, they do provide the benefits of NRCS locally-led planning in less time and at a reduced cost. The benefits include:

- Quick and inexpensive tools for setting priorities and taking action
- Providing a level of detail that is sufficient for identifying actions that can be taken with no further watershed-level studies or analyses
- Actions to be taken may require further Federal or State permits or ESA or NEPA analysis but these activities are part of standard requirements for use of best management practices (BMPs) and conservation systems
- Identifying where further detailed analyses or watershed studies are needed
- Plans address multiple objectives and concerns of landowners and communities
- Plans are based on established partnerships at the local and state levels
- Plans enable landowners and communities to decide on the best mix of NRCS programs that will meet their goals
- Plans include the full array of conservation program tools (i.e. cost-share practices, easements, technical assistance)


Rapid Watershed Assessments provide information that helps land-owners and local leaders set conservation priorities. The Frenchman Watershed is located in the Republican River Basin, on the north eastern plains of Colorado. The watershed covers 1,055,709 acres in Colorado and Nebraska, with 545,452 acres in CO. Approximately 584 farms and ranches covering 861,205 acres in the entire watershed. As of April 2005, there are 17,767 acres of land in the Conservation Reserve Program.






COLORADO County	County Acres	County Acres in FRENCHMAN Watershed	% of County in the Wa- tershed	% of Watershed in the County
Logan	1,180,481	168,575	14.3%	16.0%
Phillips	440,331	348,349	79.1%	33.0%
Sedgwick	351,884	94	0.03%	0.01%
Yuma	1,516,787	28,434	1.9%	2.7%
NEBRASKA				
Chase	575,123	311,876	54.2%	29.5%
Dundy	589,983	35,335	6.0%	3.3%
Hayes	457,410	56,411	12.3%	5.3%
Hitchcock	460,794	106,635	23.1%	10.1%
		1,055,708		

Frenchman Watershed - 10250005



Satellite Imagery: arc IMS Server - Geography Network Services hosted by ESRI




Common Resource Areas (CRA): Geographical areas where resource concerns, problems, and treatment needs are similar. Landscape conditions, soil, climate, human considerations, and other natural resource information are used to determine the geographical boundaries of the common resource area.

MLRA	CRA	CRA NAME	CRA DESCRIPTION
72	72.1	Central High Tableland	The Central High Tableland CRA is broad, level to gently rolling, loess mantled tableland. Local relief is measured in feet on the tableland tens of feet and major river valleys bordered by steep slopes. Soils are deep. Presettlement vegetation was short grass prairies. Nearly all of this area in cropland, both dryland small grain crops and irrigated corn and grain sorghum.

Physical Description


This area is characterized by broad, undulating to rolling plains dissected by streams and rivers. The highest elevations are on the northwestern side of the watershed and gently slopes down to the lowest elevation to the southeast.


The vast majority of the Frenchman Watershed consists of irrigated and dryland crops.



Land Ownership

Approximately 523,336 acres in the Colorado portion of the Frenchman Watershed are privately owned. There are 22,094 acres of state controlled land and no federally controlled lands.






EDENI HIMANI I OLOFAGO LADO			
FRENCHMAN COIORAGO LANG Use	Total Acreage	Vegetation	Acreage
		Agriculture Land	48.2
Cropland	417,245	Dryland Ag	313,058.8
		Irrigated Ag	104,137.8
		Grass Dominated	61,950.6
		Grass/Forb Mix	9.4
Rangeland/Grassland	124,006	Grass/Yucca Mix	131.5
		Sagebrush/Grass Mix Soil	60,256.5
			1,658.3 2,414.3
Riparian	3,409	Herbaceous Riparian Riparian	2,414.3 995.1
Kiparian	3,407	Shrub Riparian	87.8
Water	336.6	Water	336.6
Other	843	No Data	6.3
Other	043	Urban/Built Up	836.3
Total Colorado Watershed Acre	es		545,927
FRENCHMAN Nebraska Land Use	Total Acreage	Vegetation	Acreage
		Pasture/Hay	16,748.7
Cropland	265,297	Row Crops	173,988.7
Cropiana	203,277	Small Grains	69,984.2
		Fallow	4,575.5
Rangeland/Grassland	240,257	Grasslands/Herbaceous	240,256.5
Forest	467	Deciduous Forest	297.5
		Evergreen Forest	169.0 6.7
Riparian	893	Woody Wetlands Emergent Herbaceous Wetlands	886.5
Water	1,402	Water	1,401.8
		Bare Rock/Sand/Clay	345.0
Other	786	Low Intensity Residential	395.3
Other	700	Commercial/Industrial/Transportation	293.7
		Urban/Recreational Grasses	96.8

Precipitation

Droughts are common in the watershed as with the rest of Colorado. Statewide ,in the 1900's alone, four prolonged dry spells occurred. There was one in the 1910s. Another, in the '30s, caused the dust-bowl period. The second worst drought on record in the state occurred in the mid-50s. A series of hot, dry summers following a period of scant mountain snowpack created water shortages. The fourth drought hit parts of Colorado in the late 1970s. In this century, the most severe drought since 1723 hit the state in 2002. Prior to the 1700's, researchers looking at tree ring records have found evidence of even more severe droughts, some lasting many years.




Rainfall occurs as frontal storms in the spring and early summer and high intensity, convective thunderstorms in late summer. Maximum precipitation occurs from mid spring through late autumn. Precipitation in winter is snow. The average annual temperature is from 36 to 65 degrees F, and the frost free period averages 147 days.




Ecological Sites

The plant community on an ecological site is typified by an association of species that differs from that of other ecological sites in the kind and/or proportion of species or in total production.

Ecological Site maps give an overall indication of the soils plant relationship in the area. More detailed descriptions of ecological sites are provided in the Field Office Technical Guide (FOTG). The FOTG is available in local offices of the Natural Resources Conservation Service (NRCS) and online at http://www.nrcs.usda.gov/technical/efotg/.



<u>Land Capability Classification</u> shows, in a general way, the suitability of soils for most kinds of field crops. Crops that require special management are excluded. The soils are grouped according to their limitations for field crops, the risk of damage if they are used for crops, and the way they respond to management. The criteria used in grouping the soils do not include major and generally expensive landforming that would change slope, depth, or other characteristics of the soils, nor do they include possible but unlikely major reclamation projects. Capability classification is not a substitute for interpretations that show suitability and limitations of groups of soils for rangeland, for woodland, and for engineering purposes.



Land Capability Classes

Class 1 - soils have few limitations that restrict their use.

Class 2 - soils have moderate limitations that reduce the choice of plants or that require moderate conservation practices.

Class 3 - soils have severe limitations that reduce the choice of plants or that require special conservation practices, or both.


Class 4 - soils have very severe limitations that reduce the choice of plants or that require very careful management, or both.

Class 5 - soils are subject to little or no erosion but have other limitations, impractical to remove, that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.


Class 6 - soils have severe limitations that make them generally unsuitable for cultivation and that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

Class 7 - soils have very severe limitations that make them unsuitable for cultivation and that restrict their use mainly to grazing, forestland, or wildlife habitat.

Class 8 - soils and miscellaneous areas have limitations that preclude commercial plant production and that restrict their use to recreational purposes, wildlife habitat, watershed, or aesthetic purposes.



Farmland Classification




Colorado had approximately 1,696,800 acres of nonfederal prime farmland recorded in 1997. This represents over 2 percent of the states total land area or 4 percent of the nonfederal land in Colorado. Nationally. 64 percent of soils classified as prime farmland are being used for cropland. In Colorado, 93 percent of the soils classified as prime farmland are being utilized as cropland.

The Wind Erodibility Index (WEI): numerical value indicating the susceptibility of soil to wind erosion, or the tons per acre per year that can be expected to be lost to wind erosion if it is assumed there is no vegetative cover or management.

Soils with an erodibility index equal to or greater than 8 are considered highly erodible.


As shown on the Wind Erodibility Index map below, most cropland soils in the Frenchman Watershed are considered highly erodible.



Surface Water Quality

Section 303(d) of the Clean Water Act requires states to identify and list all water bodies where state water quality standards are not being met. Thereafter, TMDLs compromising quantitative objectives and strategies have been or will be developed for these impaired waters within the watershed in order to achieve their water quality standards.

Surface water quality in the Frenchman Watershed is generally good. Section 303(d) of the Clean Water Act requires states to identify and list all water bodies where state water quality standards are not being met for designated uses. As indicated in the map, there are no 303(d) listed streams in the watershed. The Frenchman River is designated as Primary Contact Recreation, Aquatic Life Warm I, and Agriculture. Updates to the 303d/TMDL list can be found at: http://www.cdphe.state.co.us/op/wqcc/SpecialTopics/303(d)/303dtmdlpro.html



Ground Water

The High Plains Aquifer underlies the Frenchman watershed, and is the primary source of irrigation and domestic water for the area. The High Plains aquifer is an extensive regional aquifer that underlies the Great Plains states extending from South Dakota on the north to Texas and New Mexico on the south. The Denver Basin Aquifer underlies the easternmost tip of the watershed.

Ground water quality is generally good, although moderately to very hard. Total dissolved solids in the aquifer have risen significantly since the early 1900s, and in some areas, the water may exceed drinking water standards for sulfate, chloride, fluoride, iron and arsenic. These concentrations may be naturally derived from geologic sources.

Era	System	Series	Strati- graphic Unit	Unit Thickness (feet)	Physical Characteristics	Hydro- geologic Unit	Hydrologic Characteristics
		Holocene and	Valley-fill deposits	0 to 60	Stream deposits of gravel, sand, silt, clay associated with the most recent cycle of erosion and deposition along present streams		Shallow water-table aquifer(s). Well yields range from 500 to more than 1,000 gpm in several river valleys
	Pleistoce		Dune sand	0 to 300	Fine to medium sand with small amounts of clay, silt, and coarse sand formed into hills and ridges by the wind		Typically lies above the water table; has a high infiltration rate and is important for ground-water recharge
	Quaternary Pleistor Tertiary		Loess	0 to 250	Silt with lesser amounts of very fine sand and clay deposited as windblown dust	High	Lies above the water table and does not yield water; serves for minor recharge
oic		Pleistocene	Unconsolidated alluvial deposits	0 to 550	Stream deposits of gravel, sand, silt, and clay locally cemented by calcium carbonate into caliche or mortar beds	Plains aquifer	Primary portion of the High Plains aquifer; mostly unconfined; yields
Cenoz		M:	Ogallala Formation	0 to 700	Poorly sorted clay, silt, sand, and gravel generally uncon- solidated; forms caliche layers or mortar beds when cemented by calcium carbonate; Ogallala makes up large part of High Plains aquifer		range from 100 to 3,100 gpm; typi- cally less than 300 gpm in Colorado; Ogallala is the most significant High Plains aquifer resource
		Pilocene	Arikaree Group	0 to 1,000	Predominantly massive, very-fine to fine-grained sand- stone with localized beds of volcanic ash, silty sand, silt- stone, claystone, sandy clay, limestone, marl, and mortar beds; part of the High Plains aquifer		Can be confined; moderately permeable. May yield up to 200 gpm in localized areas
		Oligocene	White River Group	0 to 700	Upper unit, Brule Formation, is considered part of the High Plains aquifer in Colorado, predominantly massive sand- stone containing sandstone beds and channel deposits) 	Typically confined, except at outcrop; yields typically less than 100 gpm
L			Огоир		Lower unit, Chadron Formation, mainly consists of varicolored, bentonitic, loosely to moderately cemented clay and silt		Chadron is mostly impermeable

From Gutentag and others, 1984

Geology

The Frenchman River lies within the Ogallala formation, and dips into Cretaceous Pierre shale on the eastern edge of the watershed. Eolian sands and silt cover much of the uplands surrounding the river.



Threatened & Endangered Species
State & Federally Threatened, Endangered & Candidate Species as well as Species of Special Concern in Frenchman Watershed

tershed									
	Common Name	Scientific Name	Class	Federal Status	State Status	Comments			
	Bald Eagle	Haliaeetus leucocephalus	Birds	None	Threatened	May migrate through water- shed			
TEN A	Black-footed Fer- ret	Mustela ni- gripes	Mammals	Endangered	Endangered	No current records of oc-currence			
	Black-tailed Prai- rie Dog	Cynomys ludo- vicianus	Mammals	None	Concern	Occurs in the watershed			
Ž	Burrowing Owl	Athene cunicu- laria	Birds	None	Threatened	Occurs in the watershed			
	Common Garter Snake	Thamnophis sirtalis	Reptiles	None	Concern	May occur in the watershed			
	Cylindrical paper- shell	Anodontoides ferussacianus	Gastro- pods	None	Concern	May occur in the watershed			
A C	Ferruginous Hawk	Buteo regalis	Birds	None	Concern	Occurs in the watershed			
	Long-Billed Cur- lew	Numenius americanus	Birds	None	Concern	Occurs in the watershed			
	Mountain Plover	Charadrius montanus	Birds	None	Concern	Occurs in the watershed			
S -3	Northern Cricket Frog	Acris crepitans	Amphibi- ans	None	Concern	May occur in the watershed			
	Northern leopard frog	Rana pipiens	Amphibi- ans	None	Concern	Occurs in the watershed			
55	Plains Leopard Frog	Rana blairi	Amphibi- ans	None	Concern	Occurs in the watershed			

Threatened & Endangered Species continued

State & Federally Threatened, Endangered & Candidate Species as well as Species of Special Concern in Frenchman Watershed


	Common Name		Scientific Class		State Status	Comments
	Plains Sharp- tailed Grouse	Tympanuchus phasianellus jamesii	Birds	None	Endangered	Occurs in the wa- tershed
	Swift fox	Vulpes velox	Mammals	None	Concern	Occurs in the wa- tershed
K	Yellow mud turtle	Kinosternon flavescens	Reptiles	None	Concern	May occur in the watershed

Shortgrass prairie, sandsage-mixed grass rangeland, and both irrigated and dry cropland are the dominant terrestrial habitat types in this watershed. Burrowing owl, mountain plover, black-tailed prairie dog, and swift fox are representative species for the shortgrass habitat. Greater prairie chickens use the sand sage-mixed grass rangeland habitats. Water is scarce and the native species in this watershed are those that can survive without abundant water supplies. Riparian areas, playa lakes, and the occasional stock pond provide seasonal to intermittent aquatic habitats. Economically important wildlife species that occur in large areas of the watershed include mule and white-tailed deer, mourning dove, and pheasant. Greater prairie chicken occur in the northwestern and southeastern parts of the watershed. Plains sharp-tailed grouse are known in the northwestern part of the watershed. Bobwhite quail, wild turkey, and snow geese occur in the Frenchman Creek riparian area. Pronghorn (antelope) may be found in the southeastern corner of the watershed.

Social Data

Social Data	Logan	Phillips	Sedwick	Yuma
	9			
Demographics (US Census, American Factfinder)				
Total population		4,480	2,747	9,841
Male		2,164	1,374	4,840
Female		2,316	1,373	5,001
Median age (years)		39.8	43.2	37.3
White		4,168	2,486	9,267
Black or African American		9	14	11
American Indian and Alaska Native		13	4	28
Asian		18	21	7
Native Hawaiian and Other Pacific Islander		1	2	2
Some other race		211	164	407
Hispanic or Latino (of any race)		527	314	1268
Economic Characteristics (US Census, American Fact	tfinder)			
In labor force (population 16 years and over)		2,039	1,340	4,919
Median household income (dollars)		32,177	28,278	33,169
Median family income (dollars)		38,144	33,953	39,814
Per capita income (dollars)		16,394	16,125	16,005
Families below poverty level		110	62	235
Individuals below poverty level		507	270	1244
County Agricultural Characteristics (Colorado Agricu	ıltural Census, cou	nty data tables)		
Farms (number)	930	334	188	864
Land in farms/ranches (acres)	1,111,135	470,837	274,243	1,351,010
Average size farm/ranch (acres)	1,195	1,410	1,459	1,567
Median size farm (acres)	608	1,000	830	1,000
Average age of farmer or rancher	52.8	53.4	56.4	52.7
Net cash return from ag sales (\$1,000)	5,092	13,313	7,716	58,023
Cattle and calves (number)	185,000	30,000	15,000	250,000

Watershed Natural Resource Concerns



Arikaree Conservation District Priorities:

The Colorado Conservation Districts identified and prioritized these resource concerns during facilitated public meetings and they are included in their Long Range Plans. Higher scores indicate higher priority.

Resource Concern By Priority	Haxtun	South Platte	Yuma	Total
Water Quality	4	5	6	15
Soil Erosion	6	6		12
Water Quantity	5		6	11
Invasive Weeds	3	4		7
Tree Planting	2	3		5
Soil Quality			4	4
Grazing Management			3	3
Wildlife Habitat	2			2
Public Relations	1			1

Selected Conservation Application Data				Frenchman 10250005			
	FY 2002	FY 2003	FY 2004	FY 2005	FY 2006	FY 2007	Total
Total Conservation Systems Planned (Acres)	36,781	18,775	Not Available	17,442	11,286	8,720	93,004
Total Conservation Systems Applied (Acres)	22,950	23,249	Not Available	22,481	15,494	28,177	112,351
Practices							
Prescribed Grazing	4,104	354	357	7,536	2,427	1,598	16,376
Upland Wildlife Habitat Management	1,356	3,668	1,127	1,355	670	7,208	15,384
Conservation Cropping System	Not Available	Not Available	3,975	3,925	5,405	12,436	25,741
Residue Management	18,281	3,606	4,529	8,076	1,865	7,361	43,718
Irrigation Water Management	5,255	6,592	2,049	1,236	2,124	5,760	23,016

Conservation Systems to Address Major Resource Concerns

Primary Resource Concern:	Rangeland	d Health			
Conservation System Description:	adequate proper sto	recovery oppor ocking of animal	ned management to tunity between gr s. Estimate 107,6 ed ranches of 3,00	Based on Conservation System Guide Code: CO 72.1-GR-01-R-Grazing	
Practices		Unit	Quantity	Cost/Unit (\$)	Estimated Cost per Median Sized Ranch (\$)
Prescribed Grazing					
Fence (382)		Ft.	15,000	0.6	9,000
Pest Management (595)		Ac.	300	15	4,500
Pipeline (516)		Ft.	12,000	2.4	28,800
Upland Wildlife Habitat Management (645)		Ac.	300	na	0
Watering Facility (614)		No.	4	410	1,640
Windbreak/Shelterbelt Establishment (380)		Ft.	1,000	.85	850
Costs to apply prescribed grazing p median sized ranch of 3,000 acres	er	No.	36		85,290
				Subt	otal Rangeland costs: \$3,070,440

Conservation Systems to Address Major Resource Concerns (cont'd)

Primary Resource Concern:	Water Quality								
Conservation System Description:	Upgrading Sprink and Pest Mgt.	kler irrigatior	Reference Conservation System Guide Code: CO 72.1-CR-Sprinkler-R-2						
Practices		Unit	Quantity	Cost/Unit (\$)	Estimated Cost (\$)				
Irrigation Water Management (4	49)*	Ac	65,000	10.20	663,000				
* includes re-bowl, renozzle, and	* includes re-bowl, renozzle, and IWM								
Nutrient Management (590)		Ac	70,000	5	350,000				
Pest Management (595)		Ac	70,000	15	\$1,050,000				

Primary Resource Concern:	Soil Erosion By Wind on dryland crops								
Conservation System Description:	Seasonal residue and Pest Mgt	Seasonal residue management with Conservation crop rotation, Nutrient and Pest Mgt							
Practices		Unit	Quantity	Cost/Unit (\$)	Estimated Cost (\$)				
Conservation Crop Rotation (328)		Ac	70,470	5	352,350				
Residue Mgmt, Seasonal (344)		Ac	70,470	5	352,350				
Nutrient Management (590)		Ac	20,000	5	100,000				
Pest Management (595)		Ac	20,000	15	300,000				
Subtotal Costs Dryland Crops: \$1,140,700									

General Effects, Impacts, and Estimated Costs of Application of Conservation Systems				
Landuse	Resource Concern	Measurable Effects	Non-measurable Effects	Estimated Cost (\$)
Rangeland	Plants		Improved plant condition, productivity, health and vigor. Grazing animals have adequate feed, forage, and shelter. Wildlife habitat is sustained or improved.	3,070,440
Dryland Crop	Soil	88,200 Total Tons/Year saved	Cropland sustainability	1,140,700
Irrigated Crop	Water		Nutrients and organics are stored, handled, disposed of, and managed so that surface water uses are not adversely affected.	1,050,000
	L	Es	timated Total Costs to Address Major Resource Co	ncerns: \$5,261,140

FOOTNOTES/BIBLIOGRAPHY

303(d) listed streams within the Watershed were created using data from Colorado Department of Public Health & Environments' Water Quality & Control Commission. Impaired streams are current as of April 30, 2006. For a list of all Colorado impaired streams, locations and priority ratings, visit http://www.cdphe.state.co.us/regulations/wqccregs/100293wqlimitedsegtmdls.pdf. Stream data from National Hydrologic Dataset http://nhd.usgs.gov

Threatened and Endangered Species information was gathered using data from the Colorado Division of Wildlife (CDOW) Natural Diversity Information Source (NDIS). NDIS GIS data may be downloaded at http:// <a href="http:// <a href="http:// <a href="http:// and is.nrel.colostate.edu">http:// <a href="http:// http://wildlife.state.co.us/Wildlife.state.co.us/Wildlife.species/SpeciesOfConcern/ThreatenedEndangeredList/ http://mountainprairie.fws.gov/endspp/CountyLists/COLORADO.htm

Resource Concerns were identified using the Colorado Association of Conservation Districts' (CACD) long range (10 year) plans from the period of 1996-2000. Only the top three environmental resource concerns for each district were used. For more information on Colorado's Conservation Districts, visit http://www.cacd.us.

Maps were generated using Soil Survey Geographic Database (SSURGO) tabular and spatial data. SSURGO data was downloaded for the following Colorado & Nebraska surveys:

San Miguel Area (CO075) Published 01/10/2007 Chase C Phillips County (CO095) Published 01/30/2008 Dundy C Yuma County (CO125) Published 08/07/2007 Hayes C

Chase County (NE029) Published 01/17/2007 Dundy County (NE057) Published 01/17/2007 Hayes County (NE085) Published 11/21/2006 Hitchcock County (NE087) Published 11/21/2006

Vegetation data was generated using the Colorado Division of Wildlife's "Colorado Vegetation Classification Project" (CVCP) data. Completed in 2003, the CVCP is a landscape level vegetation dataset created using Landsat TM imagery and then formatted for GIS use. The species identified are an overview of the most common species associated in each cover type, in order of greatest occurrence. For more information on the Colorado Vegetation Classification Project, visit http://ndis.nrel.colostate.edu/coveg.

All border state (if applicable) vegetation data courtesy of the National Land Cover Dataset (NLCD). For more information visit http://www.mrlc.gov/mrlc2k_nlcd.asp

Common Resource Area (CRA), a subdivision of the Major Land Resource Area (MLRA), is a geographical area where resource concerns, problems, or treatment needs are similar. Geographic boundaries of a CRA are determined by landscape conditions, soil, climate, human considerations and other natural resource information. For more information on Common Resource Areas visit http://soils.usda.gov/survey/geography/cra.html.

Average Annual Precipitation data was developed through a partnership between the Natural Resources Conservation Service's (NRCS) National Water and Climate Center (NWCC), the National Cartography and Geospatial Center (NCGC), and the PRISM (the Parameter-elevation Regressions on Independent Slopes Model) group at Oregon State University (OSU), developers of PRISM. Mean annual precipitation maps were developed calculating averages of rainfall for the period of 1961-1990. For more information on PRISM data visit http://www.ncgc.nrcs.usda.gov/products/datasets/climate/docs/fact-sheet.html or for more information about technical aspects of PRISM, visit the PRISM website at http://www.ocs.orst.edu/prism.

Land Ownership (status,07/22/2006 dataset) data was obtained from the Bureau of Land Management, Colorado State Office. For more information, visit http://www.blm.gov/co/st/en/BLM_Programs/geographical_sciences/gis.html

Relief & Elevation maps were created using the National Elevation Dataset (NED), 30m Digital Elevation Model (DEM) raster product assembled by the U.S. Geological Survey (USGS). A hillshade grid was created from the 30m DEM to create a 3D effect. For more information about the NED visit http://ned.usgs.gov. The data was downloaded from the NRCS Geospatial Data Gateway at http://datagateway.nrcs.usda.gov.